


Free Executive Summary



Review of Acute Human-Toxicity Estimates for Selected Chemical-Warfare Agents

Committee on Toxicology, National Research Council

ISBN: 0-309-05749-3, 100 pages, 6 x 9, paperback (1997)

This free executive summary is provided by the National Academies as part of our mission to educate the world on issues of science, engineering, and health. If you are interested in reading the full book, please visit us online at <http://www.nap.edu/catalog/5825.html>. You may browse and search the full, authoritative version for free; you may also purchase a print or electronic version of the book. If you have questions or just want more information about the books published by the National Academies Press, please contact our customer service department toll-free at 888-624-8373.

"...recommends development of more scientifically sound toxicity values for various nerve agents such as sarin, the poison gas used in the Tokyo subway terrorist attack." National Research Council

This executive summary plus thousands more available at www.nap.edu.

Copyright © National Academy of Sciences. Permission is granted for this material to be shared for noncommercial, educational purposes, provided that this notice appears on the reproduced materials, the Web address of the online, full authoritative version is retained, and copies are not altered. To disseminate otherwise or to republish requires written permission from the National Academies Press.

Summary

No reliable acute-exposure¹ standards have been established for the particular purpose of protecting soldiers from toxic exposures to chemical-warfare (CW) agents. Some human-toxicity estimates are available for the most common CW agents—organophosphorus nerve agents and vesicants; however, most of those estimates were developed for offensive purposes (that is, to kill or incapacitate the enemy) and were intended to be interim values only.

The U.S. Army's original purpose for developing human-toxicity estimates for CW agents was to enable it to predict the number of casualties that would occur during an offensive action in which the goal was to kill or incapacitate a certain fraction of the enemy forces (for example, killing or incapacitating a minimum of 50% of the least-sensitive (most-resistant) individuals). Such an approach would actually result in more than half of the exposed individuals dying (the "bonus effect"), because a certain percentage of those exposed would be expected to be more susceptible than the least-sensitive individual. Thus, exposure under the Army's original estimates would result in substantial "over-kill." These estimates understate the toxicity of the agents and therefore are inappropriate for protecting soldiers.

¹ A one-time Copyright © National Academy of Sciences. All rights reserved.
This executive summary plus thousands more available at <http://www.nap.edu>

Because of the possibility of a chemical attack by a foreign power, the Army's Office of the Surgeon General asked the Army's Chemical Defense Equipment Process Action Team (CDEPAT) to review the toxicity data for the nerve agents GA (tabun), GB (sarin), GD (soman), GF, and VX, and the vesicant agent sulfur mustard (HD) and to establish a set of exposure limits that would be useful in protecting soldiers from toxic exposures to those agents. In the 1994 report entitled *Review of Existing Toxicity Data and Human Estimates for Selected Chemical Agents and Recommended Human Toxicity Estimates Appropriate for Defending the Soldier*, the team concluded that some of the existing human-toxicity estimates are too high and are inappropriate for use in protecting soldiers. In those cases, CDEPAT proposed new estimates for various routes of exposure—percutaneous vapor, vapor inhalation, and percutaneous liquid exposures. The proposed human-toxicity estimates are only for healthy male military personnel. They must not be used for civilians.

Before making a decision on acceptance of the human-toxicity estimates proposed by CDEPAT, the Department of the Army requested that the National Research Council (NRC) independently review the CDEPAT report to determine the scientific validity of the proposed estimates. The NRC assigned this project to the Committee on Toxicology (COT) of the Board on Environmental Studies and Toxicology. The COT convened the Subcommittee on Toxicity Values for Selected Nerve and Vesicant Agents, which prepared this report. Members of the subcommittee were selected for their recognized expertise in the fields of toxicology, medicine, pathology, biostatistics, and risk assessment. The subcommittee was charged to review the Army's proposed human-toxicity estimates for GA, GB, GD, GF, VX, and HD. Specifically, the subcommittee was charged with the following tasks:

1. Review the scientific protocols and quality of the toxicity data used in revising the human-toxicity estimates for acute exposures.
2. Review the toxicity estimates for mild and nonsevere effects and for severe and lethal effects.
3. Review the methods used in deriving the human-toxicity estimates for acute exposures.
4. Determine the appropriateness of the assumptions made in deriving the human-toxicity estimates for acute exposures.

The subcommittee was not asked to recommend new toxicity estimates

or to address the policy or operational consequences of lowering the proposed human-toxicity estimates. The subcommittee's evaluations of CDEPAT's proposed estimates for GA, GB, GD, GF, VX, and HD are summarized in Tables I through 6.

The subcommittee's conclusions concerning the scientific validity of the proposed CDEPAT estimates are grouped in four categories: (1) some estimates were judged to be scientifically valid; (2) other estimates were judged adequate to serve as interim estimates until further research is conducted; (3) some estimates need to be lowered; and (4) a few estimates need to be raised.

The toxicity data that CDEPAT used to derive its proposed estimates were generated primarily from a data base developed from the 1930s to the 1960s. The existing human-toxicity estimates were based on experiments performed 30–40 years ago using various animal species in often poorly controlled studies with vastly different protocols. In reviewing the available toxicity data for the six CW agents, the subcommittee recognized that the quality of the relevant toxicity data is marginal, but it also recognized that the Army needs "best estimates" to protect its troops from exposure. For each chemical agent, data were available for only a few adverse health effects, such as death, incapacitation, cholinesterase (ChE) inhibition, miosis (a decrease in pupil size), and rhinorrhea (running nose), vesication, and erythema. Thus, even though the subcommittee concluded that some of CDEPAT's proposed estimates are scientifically valid, those conclusions are based on a limited toxicity data base. By current standards of toxicology, the toxicity data base for the agents is inadequate, and such inadequacy is a major obstacle to the Army in developing human-toxicity estimates with statistical confidence and in developing risk-management strategies.

The subcommittee recommends that the Army convene an expert panel to develop a research strategy for deriving more scientifically sound toxicity values for the agents of concern. The panel should first consider the use of such techniques as structure-activity relationships, the uncertainty factors, and in vitro systems for estimating human-toxicity values for CW agents.

If these approaches do not appear to be useful, animal and human experimentation may be recommended. Although additional research is clearly desirable to provide improved confidence in existing data, such research should not be performed on laboratory animals until expert judgment documents the need on a case-by-case basis. It must be documented that the data to be obtained from laboratory animals is needed to make a significant improvement in the protection of human health.

TABLE 1 Evaluation of Human-Toxicity for GA

Toxicity Dose	Human-Toxicity Estimates for GA			Rationale for Subcommittee's Evaluation
	Route and Form of Exposure	Existing Estimates	CDEPAT's Proposed Estimates	
Toxicity Dose ^a	Percutaneous, vapor	20,000 mg- min/m ³	15,000 mg- min/m ³	Proposed estimate supported by human data
	Inhalation, vapor	135 mg- min/m ³	70 mg-min/ m ³	Because of inadequate data on GA for this route, CDEPAT derived the estimate by assuming that GA is 0.5 times as toxic as GB; approach reasonable but estimate should be lowered because of recommended lowering of LCt ₅₀ for GB for this route; further research recommended
Toxicity Dose ^b	Percutaneous, vapor	None	2,000 mg- min/m ³	ChE inhibition data used for proposing new recommendation
	Inhalation, vapor	None	50 mg-min/ m ³	CDEPAT's proposed estimate based on a study that indicated the ratio of ICt ₅₀ ^e /LC ₅₀ is 0.75; that assumption used to establish ECt ₅₀ for severe effects; the subcommittee recommends that the ECt ₅₀ estimate be lowered to correspond to the lowered estimate for LCt ₅₀ ; further research recommended
Adverse effects and	Inhalation, vapor	0.9 mg- min/m ³	0.5 mg-min/ m ³	Human data show that humans can tolerate higher exposures; further research recommended

About this PDF file: This new digital representation of the original work has been recomposed from XML files created from the original paper book, not from the original typesetting files. Page breaks are true to the original; line lengths, word breaks, heading styles, and other typesetting-specific formatting, however, cannot be retained, and some typographic errors may have been accidentally inserted. Please use the print version of this publication as the authoritative version for attribution.

Human-Toxicity Estimates for GA					
Toxicity Type	Route and Form of Exposure	Existing Estimates	CDEPAT's Proposed Estimates	Subcommittee's Evaluation of Proposed Estimates for GA	Rationale for Subcommittee's Evaluation
D ₅₀ ^c	Percutaneous, liquid	1,500 mg for 70-kg man	1,500 mg for 70-kg man	Proposed estimate should be lowered	No uncertainty factors used in lieu of limited animal data for proposed estimate; further research recommended
	Percutaneous, liquid	None	880 mg for 70-kg man	Proposed estimate should be lowered	In the absence of adequate human or animal data for this effect, CDEPAT established the estimate by assuming ID ₅₀ ^f /LD ₅₀ ratio of 0.6 to estimate ED ₅₀ ; the subcommittee recommends that the ED ₅₀ estimate be lowered to correspond to the lowered estimate for LD ₅₀ ; further research recommended

C₅₀^c: Vapor exposure that produces lethality in 50% of the exposed animals. Ct refers to the product of concentration (c) and exposure time (t). Note that Ct is not necessarily a constant.

LD₅₀^d: Percutaneous vapor exposure or inhalation vapor exposure causing a defined effect (e.g., incapacitation, severe effects, mild effects, threshold effects).

ED₅₀^e: Liquid dose causing lethality in 50% of the exposed animals.

ED₅₀^f: Liquid dose causing a defined effect in 50% of the exposed animals.

LD₅₀^g: Vapor exposure that produces incapacitation in 50% of the exposed population.

LD₅₀^h: Liquid dose causing incapacitation in 50% of the exposed population.

About this PDF file: This new digital representation of the original work has been recomposed from XML files created from the original paper book, not from the original typesetting files. Page breaks are true to the original; line lengths, word breaks, heading styles, and other typesetting-specific formatting, however, cannot be retained, and some typographic errors may have been accidentally inserted. Please use the print version of this publication as the authoritative version for attribution.

TABLE 2 Evaluation of Human-Toxicity Estimates for GB

Human-Toxicity Estimates for GB		Rationale for Subcommittee's Evaluation		
Toxicity Category	Route and Form of Exposure	Existing Estimates	CDEPAT's Proposed Estimates	Subcommittee's Evaluation of Proposed Estimates for GA
a EC ₅₀	Percutaneous, vapor	15,000 mg-min/m ³	10,000 mg-min/m ³	Proposed estimate is scientifically valid
	Inhalation, vapor	70 mg-min/m ³	35 mg-min/m ³	Proposed estimate should be lowered
b EC ₅₀	Percutaneous, vapor	None	1,200 mg-min/m ³	Proposed estimate is scientifically valid
	Inhalation, vapor	35 mg-min/m ³	25 mg-min/m ³	Proposed estimate should be lowered
c EC ₅₀	Inhalation, vapor	2 mg-min/m ³	0.5 mg-min/m ³	Proposed estimate should be raised
	Percutaneous, liquid	1,700 mg for 70-kg man	1,700 mg for 70-kg man	Low confidence in proposed estimate; proposed estimate should serve as interim value

Toxicity Type	Human-Toxicity Estimates for GB			Rationale for Subcommittee's Evaluation
	Route and Form of Exposure	Existing Estimates	CDEPAT's Proposed Estimates	
Systemic Toxicity ^d	Percutaneous, liquid	None	1,000 mg for 70-kg man	In the absence of adequate data on GB for this effect, CDEPAT ^a assumed that the ratio of ID_{50}^c/LD_{50} is 0.6 and used that to estimate the ED_{50} values; further research recommended

Agar: Vapor exposure that produces lethality in 50% of the exposed animals. Ct refers to the product of concentration (c) and exposure time (t). Note that Ct is not necessarily a constant.

Ex₅₀: Percutaneous vapor exposure causing a defined effect (e.g., incapacitation, severe effects, mild effects, threshold effects).

$^{222}\text{RnLD}_{50}$: Liquid dose causing lethality in 50% of the exposed animals.

ED₅₀: Liquid dose causing a defined effect in 50% of the exposed animals.

50: Liquid dose causing incapacitation in 50% of the exposed population.

TABLE 3 Evaluation of Human-Toxicity Estimates for GD

Toxicity Agent	Human-Toxicity Estimates for GD			Subcommittee's Evaluation of Proposed Estimates for GA	Rationale for Subcommittee's Evaluation
	Route and Form of Exposure	Existing Estimates	CDEPAT's Proposed Estimates		
GD ^a Threshold effects	Percutaneous, vapor	None	2,500 mg- min/m ³	Proposed estimate is scientifically valid	Proposed estimate based on assumption that GD is 4 times more toxic than GB for percutaneous exposure
	Inhalation, vapor	70 mg-min/ m ³	35 mg-min/ m ³	Proposed estimate should be lowered	Proposed estimate based on the assumption that GD and GB are equipotent via this route; subcommittee recommends that LC ₅₀ estimate for GD be lowered to correspond to lowered estimate for GB; further research recommended
GD ^b Threshold effects	Percutaneous, vapor	None	300 mg-min/ m ³	Proposed estimate should serve as an interim value	In the absence of adequate human or animal data, proposed estimate based on assumption that GD is 4 times more toxic than GB for percutaneous exposure; further research recommended
	Inhalation, vapor	35 mg-min/ m ³	25 mg-min/ m ³	Proposed estimate should be lowered	In the absence of adequate human or animal data, proposed estimate based on assumption that potencies of GD and GB are comparable; EC ₅₀ estimate for GD should be lowered to correspond to the lowered estimate for GB; further research recommended

About this PDF file: This new digital representation of the original work has been recomposed from XML files created from the original paper book, not from the original typesetting files. Page breaks are true to the original; line lengths, word breaks, heading styles, and other typesetting-specific formatting, however, cannot be retained, and some typographic errors may have been accidentally inserted. Please use the print version of this publication as the authoritative version for attribution.

Human-Toxicity Estimates for GD					Rationale for Subcommittee's Evaluation
Toxicity Type	Route and Form of Exposure	Existing Estimates	CDEPAT's Proposed Estimates	Subcommittee's Evaluation of Proposed Estimates for GA	
Inhalation, vapor	Inhalation, vapor	None	0.2 mg-min/m ³	Proposed estimate should be raised	In the absence of adequate human or animal data, proposed estimate based on assumption that GD is 2.5 times more potent than GB for mitotic effects; subcommittee recommends that the LCt ₅₀ estimate for GD be raised to correspond to the recommended raised estimate for GB; further research recommended
	Percutaneous, liquid	350 mg for 70-kg man	350 mg for 70-kg man	Proposed estimate should serve as an interim value	
Percutaneous, liquid	Percutaneous, liquid	None	200 mg for 70-kg man	Proposed estimate should serve as an interim value	In the absence of adequate human or animal data, proposed estimate was derived using the ID ₅₀ /LD ₅₀ ratio of 0.6; the subcommittee recommends that CDEPAT's proposed estimate serve as an interim value; further research recommended

t₅₀: Vapor exposure that produces lethality in 50% of the exposed animals. Ct refers to the product of concentration (c) and exposure time (t). Note that Ct is not necessarily a constant.

EC₅₀: Percutaneous vapor exposure or inhalation vapor exposure causing a defined effect (e.g., incapacitation, severe effects, mild effects, threshold effects).

LD₅₀: Liquid dose causing lethality in 50% of the exposed animals.

ED₅₀: Liquid dose causing a defined effect in 50% of the exposed animals.

ID₅₀: Liquid dose causing incapacitation in 50% of the exposed population.

TABLE 4 Evaluation of Human-Toxicity Estimates for GF

Human-Toxicity Estimates for GF					Rationale for Subcommittee's Evaluation
Toxicity Dose	Route and Form of Exposure	Existing Estimates	CDEPAT's Proposed Estimates	Subcommittee's Evaluation of Proposed Estimates for GA	
Toxicity Dose ^a	Percutaneous, vapor	15,000 mg- min/m ³	2,500 mg- min/m ³	Proposed estimate should serve as an interim value	Rationale for the CDEPAT estimate not supported by data; further research recommended In the absence of adequate data, proposed estimate based on assumption that GF, GD, and GB are equipotent; approach is reasonable; because LCt ₅₀ for GB was recommended to be lowered, proposed value for GF should be lowered correspondingly; further research recommended
	Inhalation, vapor	None	35 mg-min/ m ³	Proposed estimate should be lowered	
Toxicity Dose ^b	Percutaneous, vapor	None	300 mg-min/ m ³	Proposed estimate should serve as an interim value	Proposed estimate based on assumption that GF and GD are equipotent; approach is reasonable; further research recommended In the absence of adequate data, proposed estimate based on assumption that GF, GD, and GB are equipotent; approach is reasonable; because ECt ₅₀ ⁸ for severe effects for GB and GD were recommended to be lowered, proposed value for GF should be lowered correspondingly; further research recommended
	Inhalation, vapor	None	25 mg-min/ m ³	Proposed estimate should be lowered	

About this PDF file: This new digital representation of the original work has been recomposed from XML files created from the original paper book, not from the original typesetting files. Page breaks are true to the original; line lengths, word breaks, heading styles, and other typesetting-specific formatting, however, cannot be retained, and some typographic errors may have been accidentally inserted. Please use the print version of this publication as the authoritative version for attribution.

Human-Toxicity Estimates for GF					Rationale for Subcommittee's Evaluation
Toxicity Type	Route and Form of Exposure	Existing Estimates	CDEPAT's Proposed Estimates	Subcommittee's Evaluation of Proposed Estimates for GA	
Cotton dusts	Inhalation, vapor	None	0.2 mg-min/m ³	Proposed estimate should be raised	In the absence of adequate human or animal data, the proposed estimate based on assumption that GF and GD are equipotent; approach is reasonable; because EC _{I50} for mild effects for GD was recommended to be raised, proposed value for GF should be raised correspondingly; further research recommended
	Percutaneous, liquid	None	350 mg for 70-kg man	Proposed estimate should serve as an interim value	In the absence of adequate human or animal data, proposed estimate based on assumption that GF and GD are equipotent; approach is reasonable; further research recommended
	Percutaneous, liquid	None	200 mg for 70-kg man	Proposed value should serve as an interim value	In the absence of adequate human or animal data, the proposed estimate based on assumption that GF and GD are equipotent; approach is reasonable; further research recommended

EC_{I50}: Vapor exposure that produces lethality in 50% of the exposed animals. Ct refers to the product of concentration (c) and exposure time (t). Note that Ct is not necessarily a constant.

EC_{I50}: Percutaneous vapor exposure or inhalation vapor exposure causing a defined effect (e.g., incapacitation, severe effects, mild effects, threshold effects).

ED₅₀: Liquid dose causing lethality in 50% of the exposed animals.

ED₅₀: Liquid dose causing a defined effect in 50% of the exposed animals.

TABLE 5 Evaluation of Human-Toxicity Estimates for VX

Toxicity Endpoint	Human-Toxicity Estimates for VX				Rationale for Subcommittee's Evaluation
	Route and Form of Exposure	Existing Estimates	CDEPAT's Proposed Estimates	Subcommittee's Evaluation of Proposed Estimates for GA	
LD ₅₀ ^a	Percutaneous, vapor	None	150 mg-min/ m ³	Proposed estimate should be considered an interim value	Degree of confidence in data is low to moderate; further research recommended
	Inhalation, vapor	30 mg-min/ m ³	15 mg-min/ m ³	Proposed estimate should be lowered	Degree of confidence in data is low to moderate; further research recommended
NOAEL ^b	Percutaneous, vapor	None	10 mg-min/ m ³	Proposed estimate should be considered an interim value	Degree of confidence in data is low; a no-observed- adverse-effect level (NOAEL) was not defined; further research recommended
	Percutaneous, vapor	None	25 mg-min/ m ³	Proposed estimate should be considered an interim value	Degree of confidence low to moderate; further research recommended
Sublethal effects	Inhalation, vapor	25 mg-min/ m ³	10 mg-min/ m ³	Proposed estimate should be considered an interim value	Insufficient data; further research recommended
	Inhalation, vapor	0.09 mg- min/m ³	0.09 mg- min/m ³	Proposed estimate is scientifically valid	Available human data support the proposed estimate
Sublethal effects	Percutaneous, liquid	10 mg/70- kg man	5 mg/70-kg man	Proposed estimate should be lowered	Animal data indicate that the proposed estimate is too high; furthermore, no uncertainty factor used in lieu of variability associated with dermal penetration of various regions of body; further research recommended

About this PDF file: This new digital representation of the original work has been recomposed from XML files created from the original paper book, not from the original typesetting files. Page breaks are true to the original; line lengths, word breaks, heading styles, and other typesetting-specific formatting, however, cannot be retained, and some typographic errors may have been accidentally inserted. Please use the print version of this publication as the authoritative version for attribution.

Human-Toxicity Estimates for VX				
Toxicity Type	Route and Form of Exposure	Existing Estimates	CDEPAT's Proposed Estimates	Subcommittee's Evaluation of Proposed Estimates for GA
LD ₅₀ ^d Highly persistent	Percutaneous, liquid	5 mg/70-kg man	2.5 mg/70kg man	Proposed estimate should be lowered
ED ₅₀ ^e Effects				The ED ₅₀ is based on the ID ₅₀ /LD ₅₀ ratio; the subcommittee recommends that the LD ₅₀ be lowered, therefore, the ED ₅₀ should be lowered correspondingly; further research recommended

LD₅₀: Vapor exposure that produces lethality in 50% of the exposed animals. Ct refers to the product of concentration (c) and exposure time (t). Note that Ct is not necessarily a constant.

LD₅₀: Percutaneous vapor exposure or inhalation vapor exposure causing a defined effect (e.g., incapacitation, severe effects, mild effects, threshold effects).

LD₅₀: Liquid dose causing lethality in 50% of the exposed animals.

ED₅₀: Liquid dose causing a defined effect in 50% of the exposed animals.

ED₅₀: Liquid dose causing incapacitation in 50% of the exposed population.

TABLE 6 Evaluation of Human-Toxicity Estimates for HD

Toxicity Dose Threshold Effects	Human-Toxicity Estimates for HD				Rationale for Subcommittee's Evaluation
	Route and Form of Exposure	Existing Estimates	CDEPAT's Proposed Estimates	Subcommittee's Evaluation of Proposed Estimates for GA	
a 50	Percutaneous, vapor	10,000 mg min/m ³	5,000 mg- min/m ³	Proposed estimate should be lowered	Estimate might be too high because data from the most- sensitive species (rats and mice) not used; further research recommended
	Inhalation, vapor	1,500 mg min/m ³	900 mg-min/ m ³	Proposed estimate is scientifically valid	CDEPAT averaged LC ₅₀ data in several animal species; in the absence of data on humans, that approach is reasonable
b 50	Percutaneous, vapor	None	50 mg-min/ m ³ (moderate temperature); 25 mg-min/ m ³ (hot temperature)	Proposed estimates should serve as interim values	In the absence of details on studies on which estimates were based, proposed estimate should be considered interim value; further research recommended
	Percutaneous, vapor	2,000 mg- min/ m ³ (moderate temperature) 1,000 mg min/ m ³ (hot temperature)	500 mg-min/ m ³ (moderate temperature); <200 mg- min/m ³ (hot temperature)	Proposed estimates are scientifically valid	Estimates based on human studies

About this PDF file: This new digital representation of the original work has been recomposed from XML files created from the original paper book, not from the original typesetting files. Page breaks are true to the original; line lengths, word breaks, heading styles, and other typesetting-specific formatting, however, cannot be retained, and some typographic errors may have been accidentally inserted. Please use the print version of this publication as the authoritative version for attribution.

Human-Toxicity Estimates for HD

Toxicity Type	Route and Form of Exposure	Existing Estimates	CDEPAT's Proposed Estimates	Subcommittee's Evaluation of Proposed Estimates for GA	Rationale for Subcommittee's Evaluation
Inhalation, vapor	Inhalation, vapor	200 mg-min/m ³ (moderate temperature)	100 Mg-Min/m ³ moderate temperature)	Proposed estimate is scientifically valid	Proposed estimate supported by human data
Inhalation, vapor	Inhalation, vapor	>50 mg min/m ³	25 mg-min/m ³	Proposed estimate is scientifically valid	Proposed estimate supported by human data
Percutaneous, liquid	Percutaneous, liquid	7,000 mg for 70-kg man	1,400 mg for 70kg man	Proposed estimate is scientifically valid	Proposed estimate supported by a study in dogs
Percutaneous, liquid	Percutaneous, liquid	None	610 mg for 70-kg man	Proposed estimate is scientifically valid; however, it should be rounded to 600 mg for a 70-kg man to avoid appearance of precision that is not there	Proposed estimate supported by human data

LD₅₀: Vapor exposure that produces lethality in 50% of the exposed animals. Ct refers to the product of concentration (c) and exposure time (t). Note that Ct is not necessarily a constant.
LC₅₀: Percutaneous vapor exposure or inhalation vapor exposure causing a defined effect (e.g., incapacitation, severe effects, mild effects, threshold effects).
LD₅₀: Liquid dose causing lethality in 50% of the exposed animals.
ED₅₀: Liquid dose causing a defined effect in 50% of the exposed animals.

About this PDF file: This new digital representation of the original work has been recomposed from XML files created from the original paper book, not from the original typesetting files. Page breaks are true to the original; line lengths, word breaks, heading styles, and other typesetting-specific formatting, however, cannot be retained, and some typographic errors may have been accidentally inserted. Please use the print version of this publication as the authoritative version for attribution.

The experimental designs should include the following:

- Define if and when experiments with humans are appropriate.
- In the absence of human experimentation, define the most appropriate animal model for each specific toxicity value and agent, including the end points to be observed.
- Define the adequacy of the design in determining the toxicity values for healthy female as well as healthy male military personnel.
- Define the requirements for observation of reversibility of adverse health effects.
- Identify adverse health effects at the low end of the dose-response curve to determine threshold exposure levels.
- Identify confidence limits for the proposed estimates as a measure of the uncertainty of the estimated incidence of toxic effects.
- Identify potentiation or antagonistic effects from exposures to mixtures of chemical agents.
- Identify more-sensitive biological markers of exposure and effects for CW agents.

Review of Acute Human-Toxicity Estimates for Selected Chemical- Warfare Agents

Subcommittee on Toxicity Values for Selected Nerve and Vesicant
Agents

Committee on Toxicology

Board on Environmental Studies and Toxicology

Commission on Life Sciences

National Research Council

NATIONAL ACADEMY PRESS

WASHINGTON, D.C. 1997

Copyright © National Academy of Sciences. All rights reserved.

This executive summary plus thousands more available at <http://www.nap.edu>

About this PDF file: This new digital representation of the original work has been recomposed from XML files created from the original paper book, not from the original typesetting files. Page breaks are true to the original; line lengths, word breaks, heading styles, and other typesetting-specific formatting, however, cannot be retained, and some typographic errors may have been accidentally inserted. Please use the print version of this publication as the authoritative version for attribution.

National Academy Press 2101 Constitution Avenue, NW Washington, DC 20418

NOTICE: The project that is the subject of this report was approved by the Governing Board of the National Research Council, whose members are drawn from the councils of the National Academy of Sciences, the National Academy of Engineering, and the Institute of Medicine. The members of the committee responsible for the report were chosen for their special competences and with regard for appropriate balance.

This report has been reviewed by a group other than the authors according to procedures approved by a Report Review Committee consisting of members of the National Academy of Sciences, the National Academy of Engineering, and the Institute of Medicine.

The project was supported by contract DAMD 17-89-C-9086 between the National Academy of Sciences and the U.S. Department of Defense. Any opinions, findings, conclusions, or recommendations expressed in this publication are those of the author(s) and do not necessarily reflect the view of the organizations or agencies that provided support for this project.

Additional copies of this report are available from the Board on Environmental Studies and Toxicology, 2101 Constitution Ave., N.W., Washington, D.C.

Copyright 1997 by the National Academy of Sciences. All rights reserved.

Printed in the United States of America.

About this PDF file: This new digital representation of the original work has been recomposed from XML files created from the original paper book, not from the original typesetting files. Page breaks are true to the original; line lengths, word breaks, heading styles, and other typesetting-specific formatting, however, cannot be retained, and some typographic errors may have been accidentally inserted. Please use the print version of this publication as the authoritative version for attribution.

Subcommittee on Toxicity Values for Selected Nerve and vesicant Agents

- LOREN D. KOLLER** (*Chair*), Oregon State University, Corvallis, Oreg.
DONALD E. GARDNER, Inhalation Toxicology Associates, Raleigh, N.C.
DAVID W. GAYLOR, U.S. Food and Drug Administration, Jefferson, Ark.
SIDNEY GREEN, Corning Hazleton, Inc., Vienna, Va.
ROGENE F. HENDERSON, Lovelace Biomedical and Environmental Research
Institute, Albuquerque, N.Mex.
BERNARD M. WAGNER, Wagner Associates, Inc., Millburn, N.J.

Staff

- KULBIR S. BAKSHI**, Project Director
LUCY V. FUSCO, Project Assistant
RUTH E. CROSSGROVE, Editor
Sponsor.: U.S. Department of Defense

About this PDF file: This new digital representation of the original work has been recomposed from XML files created from the original paper book, not from the original typesetting files. Page breaks are true to the original; line lengths, word breaks, heading styles, and other typesetting-specific formatting, however, cannot be retained, and some typographic errors may have been accidentally inserted. Please use the print version of this publication as the authoritative version for attribution.

Committee on Toxicology

ROGENE F. HENDERSON (*Chair*), Lovelace Biomedical and Environmental Research Institute, Albuquerque, N.Mex.

DONALD E. GARDNER (*Vice-Chair*), Inhalation Toxicology Associates, Raleigh, N.C.

GERMAINE M. BUCK, State University of New York at Buffalo, N.Y.

GARY P. CARLSON, Purdue University, West Lafayette, Ind.

JACK H. DEAN, Sanofi Winthrop, Inc., Malverne, Pa.

KEVIN E. DRISCOLL, Procter & Gamble Company, Cincinnati, Ohio

ELAINE M. FAUSTMAN, University of Washington, Seattle, Wash.

CHARLES E. FEIGLEY, University of South Carolina, Columbia, S.C.

DAVID W. GAYLOR, US Food and Drug Administration, Jefferson, Ark.

JUDITH A. GRAHAM, U.S. Environmental Protection Agency, Research Triangle Park, N.C.

IAN A. GREAVES, University of Minnesota, Minneapolis, Minn.

SIDNEY GREEN, Corning Hazleton, Inc., Vienna, Va.

WILLIAM E. HALPERIN, National Institute for Occupational Safety and Health, Atlanta, Ga.

LOREN D. KOLLER, Oregon State University, Corvallis, Oreg.

DNIEL KREWSKI, Health Canada, Ottawa, Ont.

THOMAS E. MCKONE, University of California, Berkeley, Calif.

MICHELE A. MEDINSKY, Chemical Industry Institute of Toxicology, Research Triangle Park, N.C.

JOHN L. O'DONOGHUE, Eastman Kodak Company, Rochester, N.Y.

ROBERT SNYDER, Environmental and Occupational Health Sciences Institute, Piscataway, N.J.

BERNARD M. WAGNER, Wagner Associates, Inc., Millburn, N.J.

ANNETTA P. WATSON, Oak Ridge National Laboratory, Oak Ridge, Tenn.

HANSPETER R. WITSCHI, University of California, Davis, Calif.

GAROLD S. YOST, University of Utah, Salt Lake City, Utah

About this PDF file: This new digital representation of the original work has been recomposed from XML files created from the original paper book, not from the original typesetting files. Page breaks are true to the original; line lengths, word breaks, heading styles, and other typesetting-specific formatting, however, cannot be retained, and some typographic errors may have been accidentally inserted. Please use the print version of this publication as the authoritative version for attribution.

Staff

- KULBIR S. BAKSHI**, Program Director
- MARGARET E. McVEY**, Program Officer
- RUTH E. CROSSGROVE**, Editor
- CATHERINE M. KUBIK**, Senior Program Assistant
- LINDA V. LEONARD**, Senior Project Assistant
- LUCY V. FUSCO**, Project Assistant

Board on Environmental Studies and Toxicology

PAUL G. RISSE (*Chair*), Oregon State University, Corvallis, Oreg.
MAY R. BERENBAUM, University of Illinois, Urbana, Ill.
EULA BINGHAM, University of Cincinnati, Cincinnati, Ohio
PAUL BUSCH, Malcolm Pirnie, Inc., White Plains, N.Y.
EDWIN H. CLARK II, Clean Sites, Inc., Alexandria, Va.
ELLIS COWLING, North Carolina State University, Raleigh, N.C.
GEORGE P. DASTON, The Procter & Gamble Co., Cincinnati, Ohio
PETER L. DEFUR, Virginia Commonwealth University, Richmond, Va.
DAVID L. EATON, University of Washington, Seattle, Wash.
DIANA FRECKMAN, Colorado State University, Ft. Collins, Colo.
ROBERT A. FROSCH, Harvard University, Cambridge, Mass.
DANIEL KREWSKI, Health & Welfare Canada, Ottawa, Ont.
RAYMOND C. LOEHR, The University of Texas, Austin, Tex.
WARREN MUIR, Hampshire Research Institute, Alexandria, Va.
GORDON ORIANS, University of Washington, Seattle, Wash.
GEOFFREY PLACE, Hilton Head, S.C.
BURTON H. SINGER, Princeton University, Princeton, N.J.
MARGARET STRAND, Bayh, Connaughton and Malone, Washington, D.C.
BAILUS WALKER, JR., Howard University, Washington, D.C.
GERALD N. WOGAN, Massachusetts Institute of Technology, Cambridge, Mass.
TERRY F. YOSIE, Ruder Finn Inc., Washington, D.C.

About this PDF file: This new digital representation of the original work has been recomposed from XML files created from the original paper book, not from the original typesetting files. Page breaks are true to the original; line lengths, word breaks, heading styles, and other typesetting-specific formatting, however, cannot be retained, and some typographic errors may have been accidentally inserted. Please use the print version of this publication as the authoritative version for attribution.

About this PDF file: This new digital representation of the original work has been recomposed from XML files created from the original paper book, not from the original typesetting files. Page breaks are true to the original; line lengths, word breaks, heading styles, and other typesetting-specific formatting, however, cannot be retained, and some typographic errors may have been accidentally inserted. Please use the print version of this publication as the authoritative version for attribution.

Senior Staff Officers

- JAMES J. REISA**, Director
- DAVID J. POLICANSKY**, Associate Director and Program Director for Applied Ecology
- KULBIR S. BAKSHI**, Program Director for the Committee on Toxicology
- CAROL A. MACZKA**, Program Director for Toxicology and Risk Assessment
- LEE R. PAULSON**, Program Director for Information Systems and Statistics
- RAYMOND A. WASSEL**, Program Director for Environmental Sciences and Engineering

Commission on Life Sciences

- THOMAS D. POLLARD** (*Chair*), The Salk Institute, La Jolla, Calif.
FREDERICK R. ANDERSON, Cadwalader, Wickersham & Taft, Washington, D.C.
JOHN C. BAILAR III, University of Chicago, Chicago, Ill.
PAUL BERG, Stanford University School of Medicine, Stanford, Calif.
JOHN E. BURRIS, Marine Biological Laboratory, Woods Hole, Mass.
SHARON L. DUNWOODY, University of Wisconsin, Madison, Wisc.
URSULA W. GOODENOUGH, Washington University, St. Louis, Mo.
HENRY W. HEIKKINEN, University of Northern Colorado, Greeley, Colo.
HANS J. KENDE, Michigan State University, East Lansing, Mich.
SUSAN E. LEEMAN, Boston University School of Medicine, Boston, Mass.
THOMAS E. LOVEJOY, Smithsonian Institution, Washington, D.C.
DONALD R. MATTISON, University of Pittsburgh, Pittsburgh, Pa.
JOSEPH E. MURRAY, Wellesley Hills, Mass.
EDWARD E. PENHOET, Chiron Corporation, Emeryville,
EMIL A. PFITZER, Research Institute for Fragrance Materials, Hackensack, N.J.
MALCOLM C. PIKE, University of Southern California, Los Angeles, Calif.
HENRY C. PITOT III, University of Wisconsin, Madison, Wisc.
JONATHAN M. SAMET, The Johns Hopkins University, Baltimore, Md.
CHARLES F. STEVENS, The Salk Institute, La Jolla, Calif.
JOHN L. VANDEBERG, Southwest Foundation for Biomedical Research, San Antonio, Tex.
PAUL GILMAN, Executive Director

About this PDF file: This new digital representation of the original work has been composed from XML files created from the original paper book, not from the original typesetting files. Page breaks are true to the original; line lengths, word breaks, heading styles, and other typesetting-specific formatting, however, cannot be retained, and some typographic errors may have been accidentally inserted. Please use the print version of this publication as the authoritative version for attribution.

Other Recent Reports of the Board on Environmental Studies and Toxicology

Toxicologic Assessment of the Army's Zinc Cadmium Sulfide Dispersion Tests

Carcinogens and Anticarcinogens in the Human Diet: A Comparison of Naturally Occurring and Synthetic Substances (1996)

Upstream: Salmon and Society in the Pacific Northwest (1996)

Science and the Endangered Species Act (1995)

Wetlands: Characteristics and Boundaries (1995)

Biologic Markers (Urinary Toxicology (1995), Immunotoxicology (1992), Environmental Neurotoxicology (1992), Pulmonary Toxicology (1989), Reproductive Toxicology (1989))

Review of EPA's Environmental Monitoring and Assessment Program (three reports, 1994-1995)

Science and Judgment in Risk Assessment (1994)

Ranking Hazardous Waste Sites for Remedial Action (1994)

Pesticides in the Diets of Infants and Children (1993)

Issues in Risk Assessment (1993)

Setting Priorities for Land Conservation (1993)

Protecting Visibility in National Parks and Wilderness Areas (1993)

Dolphins and the Tuna Industry (1992)

Hazardous Materials on the Public Lands (1992)

Science and the National Parks (1992)

Animals as Sentinels of Environmental Health Hazards (1991)

Assessment of the U.S. Outer Continental Shelf Environmental Studies Program, Volumes I-IV (1991-1993)

Human Exposure Assessment for Airborne Pollutants (1991)

Monitoring Human Tissues for Toxic Substances (1991)

Rethinking the Ozone Problem in Urban and Regional Air Pollution (1991)

Decline of the Sea Turtles (1990)

Tracking Toxic Substances at Industrial Facilities (1990)

Copies of these reports may be ordered from the National Academy Press
(800) 624-6242 (202) 334-3313

Other Recent Reports of the Committee on Toxicology

- Toxicity of Military Smokes and Obscurants, Volume 1 (1997)
- Toxicity of Alternatives to Chlorofluorocarbons: HFC-134a and HCFC-123 (1996)
- Permissible Exposure Levels for Selected Military Fuel Vapors (1996)
- Spacecraft Maximum Allowable Concentrations for Selected Airborne Contaminants, Volume 1 (1994), Volume 2 (1996), and Volume 3 (1996)
- Nitrate and Nitrite in Drinking Water (1995)
- Guidelines for Chemical Warfare Agents in Military Field Drinking Water (1995)
- Review of the U.S. Naval Medical Research Institute's Toxicology Program (1994)
- Health Effects of Permethrin-Impregnated Army Battle-Dress Uniforms (1994)
- Health Effects of Ingested Fluoride (1993)
- Guidelines for Developing Community Emergency Exposure Levels for Hazardous Substances (1993)
- Guidelines for Developing Spacecraft Maximum Allowable Concentrations for Space Station Contaminants (1992)
- Review of the U.S. Army Environmental Hygiene Agency Toxicology Division (1991)
- Permissible Exposure Levels and Emergency Exposure Guidance Levels for Selected Airborne Contaminants (1991)

About this PDF file: This new digital representation of the original work has been recomposed from XML files created from the original paper book, not from the original typesetting files. Page breaks are true to the original; line lengths, word breaks, heading styles, and other typesetting-specific formatting, however, cannot be retained, and some typographic errors may have been accidentally inserted. Please use the print version of this publication as the authoritative version for attribution.

THE NATIONAL ACADEMIES

National Academy of Sciences
National Academy of Engineering
Institute of Medicine
National Research Council

The **National Academy of Sciences** is a private, nonprofit, self-perpetuating society of distinguished scholars engaged in scientific and engineering research, dedicated to the furtherance of science and technology and to their use for the general welfare. Upon the authority of the charter granted to it by the Congress in 1863, the Academy has a mandate that requires it to advise the federal government on scientific and technical matters. Dr. Bruce M. Alberts is president of the National Academy of Sciences.

The **National Academy of Engineering** was established in 1964, under the charter of the National Academy of Sciences, as a parallel organization of outstanding engineers. It is autonomous in its administration and in the selection of its members, sharing with the National Academy of Sciences the responsibility for advising the federal government. The National Academy of Engineering also sponsors engineering programs aimed at meeting national needs, encourages education and research, and recognizes the superior achievements of engineers. Dr. Wm. A. Wulf is president of the National Academy of Engineering.

The **Institute of Medicine** was established in 1970 by the National Academy of Sciences to secure the services of eminent members of appropriate professions in the examination of policy matters pertaining to the health of the public. The Institute acts under the responsibility given to the National Academy of Sciences by its congressional charter to be an adviser to the federal government and, upon its own initiative, to identify issues of medical care, research, and education. Dr. Harvey V. Fineberg is president of the Institute of Medicine.

The **National Research Council** was organized by the National Academy of Sciences in 1916 to associate the broad community of science and technology with the Academy's purposes of furthering knowledge and advising the federal government. Functioning in accordance with general policies determined by the Academy, the Council has become the principal operating agency of both the National Academy of Sciences and the National Academy of Engineering in providing services to the government, the public, and the scientific and engineering communities. The Council is administered jointly by both Academies and the Institute of Medicine. Dr. Bruce M. Alberts and Dr. Wm. A. Wulf are chair and vice chair, respectively, of the National Research Council.

www.national-academies.org

Preface

Due to the Existence of large stocks of chemical-warfare (CW) agents, their easy producibility from ordinary industrial chemicals, and their potential lethal effects, there is a critical need to determine as precisely as possible the exposure levels at which CW agents cause toxic effects. This information could aid in protecting soldiers in the event of a CW attack.

This report, by the Subcommittee on Toxicity Values for Selected Nerve and Vesicant Agents of the National Research Council's Committee on Toxicology, is intended to assist the U.S. Army by assessing the scientific validity of existing human-toxicity estimates for several CW agents. The estimates considered in this report were proposed recently in the Army's Chemical Defense Equipment Process Action Team (CDEPAT) report entitled *Review of Existing Toxicity Data and Human Estimates for Selected Chemical Agents and Recommended Human Toxicity Estimates Appropriate for Defending the Soldier* (1994). The report was authored by S.A. Reutter, Ph.D., and W.A. Wade, D.V.M.; it is classified "secret" and can be obtained only with permission from the director of the U.S. Army Edgewood Research, Engineering and Development Center, Edgewood, Md.

We gratefully acknowledge Carl Curling, Jerry Glasow, William

About this PDF file: This new digital representation of the original work has been recomposed from XML files created from the original paper book, not from the original typesetting files. Page breaks are true to the original; line lengths, word breaks, heading styles, and other typesetting-specific formatting, however, cannot be retained, and some typographic errors may have been accidentally inserted. Please use the print version of this publication as the authoritative version for attribution.

About this PDF file: This new digital representation of the original work has been recomposed from XML files created from the original paper book, not from the original typesetting files. Page breaks are true to the original; line lengths, word breaks, heading styles, and other typesetting-specific formatting, however, cannot be retained, and some typographic errors may have been accidentally inserted. Please use the print version of this publication as the authoritative version for attribution.

Klenke, Francis O'Donnell, Forrest Oliverson, Gerald Palmer, Sharon Reutter, Harry Salem, and Sandra Thomson (all from the U.S. Army) for providing background information. We also thank Gail Charnley (Commission on Risk Assessment and Risk Management) and Annetta Watson (Oak Ridge National Laboratory) for making presentations to the subcommittee and providing useful information.

We are grateful for the assistance of the National Research Council staff in preparing this report. Staff members who contributed to this effort are Paul Gilman, executive director of the Commission on Life Sciences; James J. Reisa, director of the Board on Environmental Studies and Toxicology; Carol A. Maczka, program director for toxicology and risk assessment; Ruth E. Crossgrove, editor; Lucy V. Fusco, project assistant, and Catherine M. Kubik, senior program assistant. We especially wish to recognize the major contributions of the project director, Kulbir S. Bakshi, who directed the preparation of the subcommittee's report. His knowledge of the scientific and technical literature and his tireless efforts to obtain information and to organize the study plan, the subcommittee meetings, and the subcommittee's report aided in the successful completion of the project.

Finally, we would like to thank all the members of the subcommittee for their dedicated efforts throughout the development of this report.

LOREN D. KOLLER, PH.D.
*CHAIR, SUBCOMMITTEE ON TOXICITY VALUES
FOR SELECTED NERVE AND VESICANT AGENTS*
ROGENE F. HENDERSON, PH.D.
CHAIR, COMMITTEE ON TOXICOLOGY

About this PDF file: This new digital representation of the original work has been recomposed from XML files created from the original paper book, not from the original typesetting files. Page breaks are true to the original; line lengths, word breaks, heading styles, and other typesetting-specific formatting, however, cannot be retained, and some typographic errors may have been accidentally inserted. Please use the print version of this publication as the authoritative version for attribution.

Contents

	Summary	1
1	Introduction and Background	17
2	Review of Acute Human-Toxicity Estimates for GA (Tabun)	20
	Percutaneous Vapor Exposure	20
	Inhalation Vapor Exposure	22
	Percutaneous Liquid Exposure	24
	Conclusions and Recommendations	25
3	Review of Acute Human-Txicity Estimates for GB (Sarin)	28
	Percutaneous Vapor Exposure	28
	Inhalation Vapor Exposure	29
	Percutaneous Liquid Exposure	31
	Conclusions and Recommendations	32
4	Review of Acute Human-toxicity Estimates for GD (Soman)	35

About this PDF file: This new digital representation of the original work has been recomposed from XML files created from the original paper book, not from the original typesetting files. Page breaks are true to the original; line lengths, word breaks, heading styles, and other typesetting-specific formatting, however, cannot be retained, and some typographic errors may have been accidentally inserted. Please use the print version of this publication as the authoritative version for attribution.

CONTENTS	xvi
Percutaneous Vapor Exposure	35
Inhalation Vapor Exposure	37
Percutaneous Liquid Exposure	38
Conclusions and Recommendations	40
5 Review of Acute Human-toxicity Estimates for GF	43
Percutaneous Vapor Exposure	43
Inhalation Vapor Exposure	45
Percutaneous Liquid Exposure	46
Conclusions and Recommendations	47
6 Review of Acute Human-toxicity Estimates for VX	50
Percutaneous Vapor Exposure	50
Inhalation Vapor Exposure	53
Percutaneous Liquid Exposure	55
Conclusions and Recommendations	56
7 Review of Acute Human-toxicity Estimates for HB	59
Percutaneous Vapor Exposure	59
Inhalation Vapor Exposure	61
Percutaneous Liquid Exposure	63
Conclusions and Recommendations	64
8 Evaluation of the Risk-estimation Procedures Used in the CDEPAT Report	67
Use of Log-Probit Analysis	67
Use of the EC ₅₀	68
Use of Confidence Limits	71
References	72
Glossary	77
Appendix	80

1—

Introduction and Background

The U.S. Army's Chemical Defense Equipment Process Action Team (CDEPAT) recently conducted an extensive review of the scientific basis for toxicity estimates in use by the Army for several chemical-warfare (CW) agents: GA, GB, GD, GF, VX, and HD. Following a detailed analysis of the toxicity of these agents and using contemporary methods of analysis, CDEPAT concluded that many of the human-toxicity estimates in use would not protect the soldier adequately (CDEPAT 1994). Recalculations of the potencies of several of the CW agents indicate that their potencies are greater than previously determined. As a result, lower exposure levels of CW agents are expected to elicit adverse effects.

Before deciding whether to implement CDEPAT's recommendations, the U.S. Department of the Army requested that the National Research Council (NRC) independently review the CDEPAT report entitled *Review of Existing Toxicity Data and Human Estimates for Selected Chemical Agents and Recommended Human Toxicity Estimates for Defending the Soldier*. The NRC assigned the project to the Committee on Toxicology (COT) of the Board on Environmental Studies and Toxicology. The COT convened the Subcommittee on Toxicity Values for Selected Nerve and Vesicant Agents, which conducted the study and prepared this report. Subcommittee members were chosen for their expertise in several specialties, including

About this PDF file: This new digital representation of the original work has been recomposed from XML files created from the original paper book, not from the original typesetting files. Page breaks are true to the original; line lengths, word breaks, heading styles, and other typesetting-specific formatting, however, cannot be retained, and some typographic errors may have been accidentally inserted. Please use the print version of this publication as the authoritative version for attribution.

toxicology, medicine, pathology, biostatistics, and risk assessment. The subcommittee was charged with determining the scientific validity of CDEPAT's proposed human-toxicity estimates for CW agents for various routes of exposure (that is, percutaneous vapor exposures, vapor inhalation exposures, and percutaneous liquid exposures). The report considers only acute¹ exposures and acute effects. It should be noted that the human-toxicity estimates for the CW agents were proposed for healthy adult male soldiers only. They must *not* be used for the general population. Specifically, the subcommittee was charged with the following tasks:

1. Review the scientific protocols and the quality of the toxicity data used in revising the human-toxicity estimates for acute exposures.
2. Review the toxicity estimates for mild and nonsevere effects and for severe and lethal effects.
3. Review the procedures used in deriving the human-toxicity estimates for acute exposures.
4. Determine the appropriateness of the assumptions made in deriving the human-toxicity estimates for acute exposures.

In reviewing the toxicity data and the proposed human-toxicity estimates for acute exposures, the subcommittee evaluated the quality of the data, the appropriateness of the procedures used in obtaining the estimates, and the assumptions made in deriving them. The subcommittee also determined whether the supporting documentation justified the proposed recommendations and whether the studies and toxicity end points were appropriate for deriving the toxicity estimates. In reviewing the proposed human-toxicity estimates, the subcommittee reviewed only the toxicity information presented in the CDEPAT report. It did not perform an independent literature search, nor did it review any data other than those presented in the report. In addition, the subcommittee was not asked to recommend new estimates or to address the policy or operational consequences of the proposed lower human-toxicity estimates.

The exposures used in the estimates are defined as follows:

- LC_{t50} is the exposure to a vapor causing lethality in 50% of a given population and is expressed as the product of air concentration (c), in

¹ A one-time exposure.